

Tension-free Midurethral Sling Surgeries for Stress Urinary Incontinence

Ming-Ping Wu, M.D.^{1,2}, Kuan-Hui Huang, M.D.³

Department of Obstetrics and Gynecology¹, Chi Mei Foundation Hospital, Tainan, Taiwan; Department of Obstetrics and Gynecology², College of Medicine, Taipei Medical University, Taipei, Taiwan; Department of Obstetrics and Gynecology³, Chang Gung Memorial Hospital, Kaohsiung, Taiwan

ABSTRACT

The surgical treatment of stress urinary incontinence (SUI) has undergone a revolutionary change since the launch of tension-free midurethral slings, including the original practices of retro-pubic and modified trans-obturator surgical approaches. Minimally invasive techniques and high success rates have led to an increasing popularity of the procedures. The short-term efficacy of trans-obturator tension-free midurethral slings is comparable to that of the retro-pubic approach. However, preliminary evidence suggests that the trans-obturator approach has a lower success rate for treatment of intrinsic sphincter deficiency (ISD). As for adverse events after different procedures, the trans-obturator route produces fewer bladder injuries and voiding difficulties, but more groin/thigh pain, vaginal injuries or erosion of the mesh as compared with the retro-pubic route. Although the short term reports on the trans-obturator approach look promising, long-term studies and randomized controlled trials (RCTs) are needed to identify the proper indications for the various types of slings and to assess efficacy and complication rates over time. **Keywords:** urinary incontinence, tension-free midurethral slings, trans-obturator vaginal tape (TOT), tension-free vaginal tape (TVT), tension-free vaginal tape- obturator system (TVT-O).

INTRODUCTION

Urinary incontinence, meaning any involuntary leakage of urine (a symptom), is defined as urethral or extra-urethral urine leakage seen during examination (a sign). Stress urinary incontinence (SUI), is the complaint of involuntary leakage upon effort or exertion (a symptom), including sneezing or coughing; it is defined clinically as the observation of involuntary leakage from the urethra synchronous with exertion or effort, sneezing or coughing; it is presumed to be due to increased abdominal pressure (a sign) [1]. With the ability to conduct more objective urodynamic evaluations, the term genuine stress incontinence (GSI) has been replaced by the preferred term urodynamic stress incontinence (USI). USI is defined through filling cystometry as an involuntary leakage of urine during increased abdominal pressure in the absence of a detrusor contraction [1]. A randomly sampled community-based study to evaluate the prevalence of urinary incontinence in a healthy population conducted by Chen et al [1] showed that a total of 53.7% of women suffered from urinary incontinence with or without related symptoms. The prevalence of stress urinary incontinence, overactive bladder, and mixed incontinence, with each category mutually

exclusive, was 18.0%, 18.6%, and 17.1%, respectively, based on patients' perceptions. The criteria of the International Continence Society (ICS) yields a lower incidence, with the prevalence of the above three conditions as 4.3%, 2.4%, and 1.8%, respectively. The occurrence of SUI increases with age (reaching 25% in the 50-65-year-old cohort). Approximately two thirds of the incontinent women reported restricting social activities because of embarrassment and inconvenience and approximately 19% reported it affected their sex life. However, only 27.1% of the women with urinary incontinence related symptoms in their study reported seeking medical services to solve these problems [2].

A brief history of anti-incontinence surgery

There have been more than 100 types of anti-incontinence surgeries invented since Kelly's sub-urethral fascia placcation reported in the year 1914 [3]. A brief summary of the history of anti-incontinence surgeries is listed in Table 1. The methods of surgical treatment for urinary incontinence vary according to the type of SUI, patients' conditions, and surgeons' preferences. Tension-free vaginal tape (TVT) (Gynecare, Ethicon Inc, Summerville, NJ), first introduced by Ulmsten in 1995 [4], has gained popularity with both gynecologists and urologists. The International Urogynecological Association (IUGA) surveyed its 152 members, including gynecologists (89%) and urologists (11%), regarding the management of pelvic floor dysfunction in 2002 and found the operations preferred for SUI were TVT (48.8%), and Burch colposuspension (44%). In SUI related to low urethral closure pressure, i.e. intrinsic sphincter deficiency (ISD), the surgery chosen was 44.6% for TVT and 32.3% for the conventional sling [5]. We have summarized the evolution of the types of anti-incontinence surgeries in Fig. 1. A review conducted in Cochrane Library in 2007 indicated that open retro-pubic colposuspension is still considered an effective treatment modality in the long run for SUI. Comparing the results of treatments in the first year, the overall continence rate achieved is approximately 85% to 90%. After five years, approximately 70% of the patients can expect to be dry. However, newer minimally invasive procedures like TVT [6] look promising in comparison to open colposuspension, but their long-term performance is not known [7]. This review will address the mechanisms, efficacy, complications and issues related to tension-free midurethral slings.

MECHANISMS AND THEORIES OF URINARY CONTINENCE

Traditionally, the pathophysiology of stress urinary incontinence emphasized the bladder neck and proximal urethral hypermobility [8]. When performing retro-pubic urethropexy for SUI, e.g. the Marshall-Marchetti-Krantz (MMK) procedure or the Burch colposuspension, the goal of the procedure is to lift the urethra up to a higher retro-pubic position [9]. Burch colposuspension attaches the endopelvic fascia at

Received: August 11, 2007 Accepted: September 27, 2007

Address correspondence to: Dr. Kuan-Hui Huang, Department of Obstetrics and Gynecology, Chang Gung Memorial Hospital, 123, Ta-Pei Road, Niao Sung Hsiang, Kaohsiung, 83305, Taiwan

E-mail: gynh2436@adm.cgmh.org.tw

the level of the bladder neck to the ipsi-lateral ileal-pectineal (Cooper) ligament, and fixes the bladder neck high in the retro-pubic space. With good results both short term (85%-90% within the first year) and long term (70% for over 5 years), it has been regarded as the gold standard of treatment [7]. Yet, the emergence of tension-free midurethral slings, designed according to integral theory [10,11] and supported by hammock theory [12], has challenged this traditional solution.

Hammock theory

The urethra lies on a supportive layer that is composed of endopelvic fascia and the anterior vaginal wall. The stability of the sub-urethral layer depends on the intact connection of the vaginal wall and endopelvic fascia to the arcus tendineus fasciae pelvis (ATFP), i.e. "white line" and levator ani muscle [12]. The increases in urethral closure pressure (the difference between bladder and urethra) during a cough arises because the urethra is compressed against a hammock-like supportive layer, rather than being truly "intra-abdominal" [12].

Integral theory

Tension-free midurethral slings have revolutionized the treatment of SUI [6,13]. The successful outcomes of the TVT procedure have been attributed not to restricting the hypermobility of the urethra and bladder neck, but to the principles of integral theory [10,11]. This theory indicates that the consequences of SUI are detachment of the pub-ourethral ligament (PUL) supporting the urethra, weakened support of the pubo-coccygeal musculature (PCM) adjacent to the urethra. During increased abdominal pressure, e.g. coughing, laughing, etc, the PCM fast twitch contraction pulls forward the upper vagina tightly around the urethra with PUL. This contraction closes the urethra off and immobilizes it while the levator plate (LP) and longitudinal muscle of the anus (LMA) pulls the bladder down and back to create a "zone of critical elasticity". Loss of elasticity here may cause the forward movement of the vagina to fail, leaving the bladder neck in the incontinent "open position" [10,11].

Dynamic kinking with "urethral knee"

Various radiologic studies demonstrating functional kinking of the midurethra during stressful maneuvers have been described. Lo et al performed ultrasound assessment of the TVT sling at 3-year follow-up,

which revealed midurethral dynamic kinking for which they coined the term "the urethral knee" [14].

DIFFERENT APPROACHES FOR TENSION-FREE MIDURETHRAL SLINGS: RETRO-PUBIC VERSUS TRANS-OBTURATOR

Tension-free midurethral sling operations include the implantation of a suburethral tape of differing materials without tension under the midurethra. It is aimed to correct the inadequate urethral support and the suburethral vaginal wall [7]. We have summarized the outcome of TVTs performed in essential institutions in various countries (Table 2).

Tension-free midurethral sling procedures can be performed using different approaches with different kinds of implants. These implants differ not only with respect to the material (polyester, polyethylene,

The evolution of anti-incontinence surgeries




Fig. 1. A brief summary of the evolution of anti-incontinence surgery. IVS: intravaginal slingoplasty; SIS: small intestine submucosa; SPARC: supra-pubic arc; TVT: tension-free vaginal tape; TVT-O: tension-free vaginal tape-obturator system.

Table 1. Brief Summary of Chronological Anti-incontinence Surgeries

Year	Author	Types of Surgeries
1914	Kelly HA [3]	Suburethral fascia plication
1917	Goeball R & Stoeckel W [56]	Using a strip of rectus fascia to encircle the urethra
1949	Marshall VF, Marchetti AA & Krantz KE [57]	Retro-pubic urethropexy (pubis periosteum)
1959	Pereyra AJ [58]	Needle-assisted transvaginal bladder neck suspension
1961	Burch JC [9]	Retro-pubic colposuspension (Cooper ligament)
1973	Stamey TA [59]	Endoscopic bladder neck suspension
1980s	Raz S [60]	Raz bladder neck suspension
1995	Ulmsten U [4]	Tension-free Vaginal Tape (TVT) (Gynecare, Ethicon Inc, Summerville, NJ)
2001	Delorme E [23]	Trans-obturator Subfascial Hammock, MONARC (American Medical System)
2003	Deval B [17]	Supra-pubic arc (SPARC) (American Medical System, Minnetonka, MN)
2003	Petros P [61]	Intra-vaginal Slingoplasty (IVS) Tunneller (Tyco Healthcare- United States Surgical, Norwalk, CT)
2003	deLeval J [24]	Trans-obturator vaginal tape (TVT-O) (Gynecare, Ethicon Inc)
2005	Mostow EN [17]	Extracellular matrix graft: absorbable sling (SIS) (Cook Biotech Inc., W. Lafayette, IN)

polypropylene, polytetrafluoroethylene, PTFE, polyethylene terephthalate, PETP, etc), but also in terms of structure (woven, knitted, monofilament and multifilament), pore size, mechanical properties, shape and surface characteristics [15]. The classification of synthetic prostheses was listed in Table 3 [16]. Type I macroporous, monofilamentous mesh is regarded in surgery as the most suitable material for stress urinary incontinence and pelvic floor reconstruction. Most commercially available tension-free midurethral slings use type I polypropylene mesh tape, with the exception of intra-vaginal slingplasty tunneller (IVS tunneller) (Tyco Healthcare-United States Surgical, Norwalk, CT), which is a type III mesh. It is also worth noting for its newly developed absorbable materials, called small intestinal submucosa (SIS) (Cook Biotech Inc., W. Lafayette, IN) [17]. SIS is a natural biomaterial harvested from the porcine small intestine and made into a biocompatible medical product using a patented process. The emergence of absorbable material may mark the beginning of a new era, however, due to limited literature on the topic, the long-term effects are still unknown.

Retro-pubic approach: top-down (TVT) or bottom-up (SPARC)

The retro-pubic approach can be done through a bottom-up vaginal approach, e.g. TVT sling (Gynecare, Ethicon Inc, Summerville, NJ); or through a top-down low abdominal approach, e.g. suprapubic arc (SPARC) sling (American Medical System, Minnetonka, MN). Two types of retro-pubic approaches are illustrated in Fig. 2. Nilsson et al reported their results after 7 years of follow-up involving 90 women. A cure rate of 81.3% was noted after a follow-up period of 91 months [13, 18]. Two randomized trials (RCTs) have reported no differences in efficacy between SPARC and TVT at 2 years. The success rates for SPARC and TVT were 83% (n=41) vs 95% (n=43), $0.05 < p < 0.1$ (12 months) [19]; and 80.7% (n=31) vs 87.1% (n=31), $p = 0.706$ (2 years) [20], neither one being significant.

Trans-obturator approach: outside-in (TOT) or inside-out (TVT-O)

TVT has been regarded as a very safe procedure. However, re-

ports from Scandinavia, Austria and other places have drawn attention to severe surgical complications relating to the penetration of the retro-pubic space [21,22]. In 2001, a trans-obturator approach was proposed for the surgical placement of sub-urethral tapes under the middle urethra, with the aim of reducing or even eliminating complications by sparing the retro-pubic space [23]. Clinical results and anatomic studies suggest this approach may be safer. The tapes pass from the thigh fold, through the obturator foramina, underneath the urethra, towards the anterior vaginal wall, without entering the pelvic region at any time during the procedure. It can be performed via an outside-in approach (TOTs), e.g. ObTape, UraTape, Monarc (American Medical System, Minnetonka, MN), etc. [23], or an inside-out approach, i.e. TVT-obturator system (TVT-O) (Gynecare, Ethicon Inc, Summerville, NJ) [24]. Two types of trans-obturator approaches are illustrated in Fig. 3. As for the efficacy for the two trans-obturator approaches, the reported results are very similar; 90% for Monarc (n=50) versus 94% for TVT-O (n=50) at 1 year follow-up, with no significant differences [25]. Although short-term data show no difference in cure rates or complications, there are no long-term studies comparing the inside-out and outside-in trans-obturator approaches [26].

Efficacy of retro-pubic versus trans-obturator approaches

According to a recent review article by Latthe et. al based on five RCTs for TVT-O versus TVT and six RCTs for TOT versus TVT, the subjective cure rates were as follows: TVT-O vs TVT (OR 0.69; 95% CI 0.42-1.15); TOT vs TVT (OR 1.04; 95% CI 0.64-1.70); total TVT-O/TOT vs TVT (OR 0.85; 95% CI 0.60-1.21), all not significant [27]. However, the trans-obturator approach is less successful in patients with ISD [28]. According to the pre-operative maximal urethral closure pressure (MUCP), the cure rates were noted as 86% if the MUCP was greater than 30 cmH₂O; compared 81% if the MUCP was 20-30 cmH₂O; and 70% if the MUCP was less than 20 cmH₂O [28]. In summary, the short-term efficacy of trans-obturator tension-free midurethral slings is comparable to the retro-pubic approach. However, preliminary evidence suggests that trans-obturator tension-free midurethral slings may have a lower success rate for treatment of ISD [27].

Table 2. Outcomes Using Tension-free Vaginal Tape (TVT) at Major Institutions in Various Countries

	Case No.	Successful	Follow-up (Mons)
Ulmsten U (Sweden) [62]	50	86%	36
Olsson U (Sweden) [63]	51	90%	36
Wang AC (Taiwan) [64]	70	84%	3-18
Jacquetin B (France) [65]	156	89.1%	12-36
Soulie M (France) [66]	120	86.7%	15.2 (6-36)
Nilsson CG (Finland) [18]	85	85%	56 (48-70)

Table 3. Classification of Synthetic Prostheses [16]

Type	Characteristics	Pore size	Brand names
Type 1	Monofilament, macroporous	> 75 μ	Gynemesh, TVT, TVT-O SPARC, Monarc, Perigee, Apogee, Prolift
Type 2	Monofilament, microporous	< 10 μ	GORE TEX
Type 3	Macroporous with multifilaments or microporous		IVS tunneller, URATAPE, SURGIPRO, MERSILENE, PARIETEX
Type 4	Submicronic	< 1 μ	




Fig. 2. Retro-pubic tension-free midurethral slings can be performed via a bottom-up vaginal approach (upper panel), e.g. TVT (A, B, C), or via a top-down low abdominal approach (lower panel), e.g. SPARC (D, E, F).




Fig. 3. The trans-obturator approach can be done outside-in with initial incisions at the groin/thigh (A, B, C), or inside-out with initial incisions at the anterior vaginal wall (D, E, F).

Proposed Indication for retro-pubic versus trans-obturator approaches

There is a paucity of data in the literature to direct physicians regarding the best indications for the use of various tension-free midurethral slings. In addition, the indications are often dependent on clinical experience and the level of comfort with the use of the various techniques. Indications for the better choice among these approaches were proposed by Silva based on common sense principles rather than rigorous study as RCTs [29].

The retro-pubic approach may be preferred over the trans-obturator approach in young, physically active patients or patients with ISD, because it helps prevent thigh/groin discomfort with exercise and the vectors of support create more urethral tension [30,31].

The trans-obturator approach may be preferred over the retro-pubic approach for obese patients, elderly patients, patients with previous retro-pubic or major abdominal surgery, or with mixed incontinence. It can make passing needles easier, reducing the risk of organ injury, reducing the risk of voiding dysfunction, and the risk of urethral obstruction [29].

Until now, there are no conclusive data to recommend the inside-out (TVT-O) over the outside-in (TOT) approach; or the bottom-up (TVT) versus the top-down approach (SPARC). Usually, surgeon preference and training plays a significant role in this decision.

COMPLICATIONS FROM TENSION-FREE MIDURETHRAL SLINGS

A nationwide analysis to evaluate the therapy-associated morbidity from TVT operations was conducted in Finland, where 1,455 patients who underwent TVT from 38 hospitals were recruited [21]. Intra-operative complications were as follows: bladder perforation; 38/1,000; active bleeding (blood loss >200 mL): 19/1,000; injury to major vessels: 0.7/1,000; nerve injury: 0.7/1,000; urethral lesions: 0.7/1,000. Post-operative complications included: retro-pubic hematoma (19/1,000); minor post-operative voiding difficulty (76/1,000); post-operative urine retention (23/1,000); post-operative urinary tract infection (41/1,000); defect in vaginal healing (7/1,000); and complications requiring laparotomy (3.4/1,000) [21]. The study concluded that the TVT procedure is a

safe method for the treatment of SUI provided that appropriate training is offered [21]. Lower urinary tract (LUT) injuries from the trans-obturator approach were reported to be 1% in a retrospective cohort study based on 390 procedures, with 241 using the outside-in and 148 using the inside-out technique. Four LUT injuries occurred; two urethral injuries (0.5%) and two bladder injuries (0.5%). All LUT injuries occurred in the outside-in group, although this difference did not reach significance ($p=0.146$) [32]. When comparing adverse events according to different approaches, the trans-obturator approach resulted in fewer bladder injuries (OR 0.12; 95% CI 0.05-0.33), and fewer voiding difficulties (OR 0.55; 95% CI 0.31-0.98); however, it was also associated with more

groin/thigh pain (OR 8.28; 95% CI 2.7-25.4), vaginal injuries or erosion of the mesh (OR 1.96; 95% CI 0.87-4.39), as compared with the retro-pubic route [27].

Urine retention and/or voiding dysfunction

Urine retention is one of the most common complications following application of tension-free midurethral slings. It is usually caused by undue tension. The symptoms include hesitancy, straining to void, incomplete emptying, urine retention, and increased post-voidal residuals [33]. It can manifest as bladder outlet obstruction, and high pressure with low flow in urodynamic studies [34]. The reported incidence ranges from 1%-17% for voiding disturbances, and




Fig. 4. Comparison of retro-pubic versus trans-obturator approach. The retro-pubic approach (A), e.g. TVT (C), SPARC (D) as the two most common commercially available kits, passes the tape through the retro-pubic space with the incision wounds located low on the abdominal wall (G). The trans-obturator approach (B), e.g. Monarc (E), TVT-O (F), passes the tape through the obturator foramen with the incision wounds located in the groin/thigh areas (H).

0%-3% for urinary retention [35]. The recommended surgical management of urine retention consists of transvaginal transection or loosening [33], urethral dilatation using Hegar dilatation [36], or lateral excision [37]. Various surgical approaches may be used to treat voiding dysfunctions following an anti-incontinence procedure. Following a vaginal or retro-pubic urethrolysis or removal of a synthetic suburethral sling, obstructive symptoms are likely to improve, symptoms of irritation may remain unchanged, and almost half will develop a recurrence of SUI. Additionally, there was a statistically significant improvement on both quality of life (QoL) questionnaires [38].

Bladder perforation

A variable and high incidence of bladder injury has been reported in the literature, ranging from 0 to 25% [6,39]. It was also reported as 0.8% (5/600) in a study by Wang [40]. There are risk factors responsible for perforation, such as previous pelvic surgery, repeated anti-incontinence surgery, esp. previous colposuspension [41]. Meanwhile, a

learning curve is also evident as regards bladder injury; the relationship between the incidence of cystotomy and the cumulative number of cases performed is inversely correlated. As the number of cases a resident completed increased, there was a slight tendency for cystotomy to decrease ($p=0.033$) [42]. Therefore, correct insertion of the trocar and intraoperative urethrocystoscopy is imperative in TVTs [40].

A prospective RCT showed that bladder injury was significantly more frequent with the retro-pubic approach compared with the trans-obturator approach (9.5 vs. 0.0%, $P=0.03$) [43]. However, unrecognized bladder injuries have also been reported with the trans-obturator approach [28]. Although intra-operative cystoscopy was not recommended for routine use, instead recommended only for selected cases in the original study design [32], some cases of inadvertent cystotomy were noted during routine cystoscopy. Therefore, some authors recommended routine cystoscopy for the trans-obturator procedure, particularly if the outside-in approach is used [44].

Sling erosion

"Erosion" is defined as the presence of foreign material within the genitourinary tract. A foreign body initially present outside the genitourinary tract and may gradually erode into the bladder, urethra, or vagina [45]. It can present as persistent vaginal discharge, partner discomfort during sexual intercourse, or even be completely asymptomatic [46]. Vaginal erosion of the TVT tape was reported as 0.9% (3/350) [47], and 1.1% (6/546) in our series [48], with incidence varying from 0.3% to 23% [49]. Sling erosions can manifest as vaginal exposure (Fig. 5C, D), or bladder exposure. The erosion rate was reported to be 2% for TVT-O, as compared with 1.5% for a TVT comparison group [50]. Concerning sling materials, complications from type I mesh, macroporous monofilament polypropylene tape (TVT) are rare 0.2%-1.2% [21]. The complication rate for ObTape (Mentor-Porges, Le Plessis Robinson, France), a non-woven polypropylene tape, is 6.1%-20% [51, 52]. With type III multifilament woven polypropylene tape, vaginal erosion occurs between 7.5%-14% of the time [53].

Most studies suggest complete removal of the eroded tape [54], but conservative treatments are available as alternative options. The exposed vaginal polypropylene slings may be managed with cautious observation. Spontaneous healing can be anticipated in some cases [46]. Sling preservation with urinary continence and patient satisfaction is a possibility for those with vaginal exposure of polypropylene mesh of less than 1 cm [46]. Vaginally exposed polypropylene mesh will become re-epithelialized within 6 weeks. If no overgrowth is evident by 3 months, excision for the eroded slings should be seriously considered [48]. Even with excision of the suburethral sling, the urethral continence function will still be effective [33,55].

CONCLUSION

The tension-free midurethral sling is a revolutionary surgical treatment for SUI. Its minimally invasive approach and success rates have led to increasing popularity of the technique. The trans-obturator approach is a potentially safer method as it spares the retro-pubic space. Although the short-term results of tension-free midurethral slings are comparable to retro-pubic colposuspension, long-term studies and RCTs are needed to




Fig. 5. Two common complications of tension-free midurethral sling surgery include bladder injury and sling erosion. Bladder perforation can either be detected (A) or undetected (B) intra-operatively in a patient with concomitant pelvic organ prolapse. Partial (C) or complete vaginal exposure (D) of sling erosions are illustrated here.

identify the proper indications for the various types of slings and to assess efficacy and complication rates over time.

REFERENCES

- Abrams P, Cardozo L, Fall M, et al: The standardisation of terminology of lower urinary tract function: Report from the Standardisation Sub-committee of the International Continence Society. *Neurourol Urodyn* 2002; **21**:167-178.
- Chen GD, Lin TL, Hu SW, Chen YC, Lin LY: Prevalence and correlation of urinary incontinence and overactive bladder in Taiwanese women. *Neurourol Urodyn* 2003; **22**:109-117.
- Kelly HA, Dumm WM: Urinary incontinence in women, without manifest injury to the bladder. 1914. *Int Urogynecol J Pelvic Floor Dysfunct* 1998; **9**:158-164.
- Ulmsten U, Petros P: Intravaginal slingplasty (IVS): An ambulatory surgical procedure for treatment of female urinary incontinence. *Scand J Urol Nephrol* 1995; **29**:75-82.
- Davila GW, Ghoniem GM, Kapoor DS, Contreras-Ortiz O: Pelvic floor dysfunction management practice patterns: A survey of members of the International Urogynecological Association. *Int Urogynecol J Pelvic Floor Dysfunct* 2002; **13**:319-325.
- Ulmsten U, Henriksson L, Johnson P, Varhos G: An ambulatory surgical procedure under local anesthesia for treatment of female urinary incontinence. *Int Urogynecol J Pelvic Floor Dysfunct* 1996; **7**:81-86.
- Lapitan MC, Cody DJ, Grant AM: Open retropubic colposuspension for urinary incontinence in women. *Cochrane Database Syst Rev* 2005; **20**:CD002912.
- Schick E, Jolivet-Tremblay M, Tessier J, Dupont C, Bertrand PE: Observations on the function of the female urethra: III: An overview with special reference to the relation between urethral hypermobility and urethral incompetence. *Neurourol Urodyn* 2004; **23**:22-26.
- Burch JC: Urethrovaginal fixation to Cooper's ligament for correction of stress incontinence, cystocele, and prolapse. *Am J Obstet Gynecol* 1961; **81**:281-290.
- Petros PE, Ulmsten UI: An integral theory and its method for the diagnosis and management of female urinary incontinence. *Scand J Urol Nephrol Suppl* 1993; **153**:1-93.
- Petros PE, Ulmsten UI: An integral theory of female urinary incontinence. Experimental and clinical considerations. *Acta Obstet Gynecol Scand Suppl* 1990; **153**:7-31.
- DeLancey JO: Structural support of the urethra as it relates to stress urinary incontinence: The hammock hypothesis. *Am J Obstet Gynecol* 1994; **170**:1713-1723.
- Nilsson CG, Falconer C, Rezapour M: Seven-year follow-up of the tension-free vaginal tape procedure for treatment of urinary incontinence. *Obstet Gynecol* 2004; **104**:1259-1262.
- Lo TS, Horng SG, Liang CC, Lee SJ, Soong YK: Ultrasound assessment of mid-urethra tape at three-year follow-up after tension-free vaginal tape procedure. *Urology* 2004; **63**:671-675.
- Cosson M, Debodinance P, Boukerrou M, et al: Mechanical properties of synthetic implants used in the repair of prolapse and urinary incontinence in women: Which is the ideal material? *Int Urogynecol J Pelvic Floor Dysfunct* 2003; **14**:169-178.
- Amid PK: Classification of biomaterials and their related complications in abdominal wall hernia. *Hernia* 1997; **1**:15-21.
- Mostow EN, Haraway GD, Dalsing M, Hodde JP, King D: Effectiveness of an extracellular matrix graft (OASIS Wound Matrix) in the treatment of chronic leg ulcers: A randomized clinical trial. *J Vasc Surg* 2005; **41**:837-843.
- Nilsson CG, Kuuva N, Falconer C, Rezapour M, Ulmsten U: Long-term results of the tension-free vaginal tape (TVT) procedure for surgical treatment of female stress urinary incontinence. *Int Urogynecol J Pelvic Floor Dysfunct* 2001; **12**(Suppl 2):S5-S8.
- Andonian S, Chen T, St-Denis B, Corcos J: Randomized clinical trial comparing suprapubic arch sling (SPARC) and tension-free vaginal tape (TVT): One-year results. *Eur Urol* 2005; **47**:537-541.
- Tseng LH, Wang AC, Lin YH, Li SJ, Ko YJ: Randomized comparison of the suprapubic arc sling procedure vs tension-free vaginal taping for stress incontinent women. *Int Urogynecol J Pelvic Floor Dysfunct* 2005; **16**:230-235.
- Kuuva N, Nilsson CG: A nationwide analysis of complications associated with the tension-free vaginal tape (TVT) procedure. *Acta Obstet Gynecol Scand* 2002; **81**:72-77.
- Tamussino K, Hanzal E, Kolle D, Ralph G, Riss P: The Austrian tension-free vaginal tape registry. *Int Urogynecol J Pelvic Floor Dysfunct* 2001; **12**(Suppl 2):S28-S29.
- Delorme E: Transobturator urethral suspension: Mini-invasive procedure in the treatment of stress urinary incontinence in women. *Prog Urol* 2001; **11**:1306-1313.
- De Leval J: Novel surgical technique for the treatment of female stress urinary incontinence: Transobturator vaginal tape inside-out. *Eur Urol* 2003; **44**:724-730.
- Debodinance P: Trans-obturator urethral sling for the surgical correction of female stress urinary incontinence: Outside-in (Monarc) versus inside-out (TVT-O) Are the two ways reassuring? *Eur J Obstet Gynecol Reprod Biol* 2007; **133**:232-238.
- Costa P, Delmas V: Trans-obturator-tape procedure--"inside out or outside in": Current concepts and evidence base. *Curr Opin Urol* 2004; **14**:313-315.
- Latthe PM, Foon R, Toozs-Hobson P: Transobturator and retropubic tape procedures in stress urinary incontinence: A systematic review and meta-analysis of effectiveness and complications. *BJOG* 2007; **114**:522-531.
- Roumeguere T, Quackels T, Bollens R, et al: Trans-obturator vaginal tape (TOT) for female stress incontinence: One year follow-up in 120 patients. *Eur Urol* 2005; **48**:805-809.
- Silva WA: Treatment of stress urinary incontinence--midurethral slings: Top-down, bottom-up, "outside-in," or "inside-out". *Clin Obstet Gynecol* 2007; **50**:362-375.
- Peschers UM, Tunn R, Buczkowski M, Perucchini D: Tension-free vaginal tape for the treatment of stress urinary incontinence. *Clin Obstet Gynecol* 2000; **43**:670-675.
- Kohli N: Tradition is yielding to new technology's advantaged, time-tested through they are not-yet. *OBS Manage* 2005; **17**:32-36.
- Abdel-Fattah M, Ramsay I, Pringle S: Lower urinary tract injuries after transobturator tape insertion by different routes: A large retrospective study. *BJOG* 2006; **113**:1377-1381.
- Klutke C, Siegel S, Carlin B, Paszkiewicz E, Kirkemo A, Klutke J: Urinary retention after tension-free vaginal tape procedure: Incidence and treatment. *Urology* 2001; **58**:697-701.
- Chassagne S, Bernier PA, Haab F, Roehrborn CG, Reisch JS, Zimmern PE: Proposed cutoff values to define bladder outlet obstruction in women. *Urology* 1998; **51**:408-411.
- Merlin T, Arnold E, Petros P, et al: A systematic review of tension-free urethropexy for stress urinary incontinence: Intravaginal slingplasty and the tension-free vaginal tape procedures. *BJU Int* 2001; **88**:871-880.
- Lo TS, Wang AC, Horng SG, Liang CC, Soong YK: Ultrasonographic and urodynamic evaluation after tension free vagina tape procedure (TVT). *Acta Obstet Gynecol Scand* 2001; **80**:65-70.
- Long CY, Lo TS, Liu CM, Hsu SC, Chang Y, Tsai EM: Lateral excision of tension-free vaginal tape for the treatment of iatrogenic urethral obstruction. *Obstet Gynecol* 2004; **104**:1270-1274.
- Segal JL, Vassallo BJ, Kleeman SD, Hungler M, Karram MM: The efficacy of the tension-free vaginal tape in the treatment of five subtypes of stress urinary incontinence. *Int Urogynecol J Pelvic Floor Dysfunct* 2006; **17**:120-124.

39. Azam U, Frazer MI, Kozman EL, Ward K, Hilton P, Rane A: The tension-free vaginal tape procedure in women with previous failed stress incontinence surgery. *J Urol* 2001; **166**:554-556.
40. Wang AC: The techniques of trocar insertion and intraoperative urethrocystoscopy in tension-free vaginal taping: An experience of 600 cases. *Acta Obstet Gynecol Scand* 2004; **83**:293-298.
41. Tamussino KF, Hanzal E, Kolle D, Ralph G, Riss PA: Tension-free vaginal tape operation: Results of the Austrian registry. *Obstet Gynecol* 2001; **98**:732-736.
42. McLennan MT, Melick CF: Bladder perforation during tension-free vaginal tape procedures: Analysis of learning curve and risk factors. *Obstet Gynecol* 2005; **106**:1000-1004.
43. David-Montefiore E, Frobert JL, Grisard-Anaf M, et al: Peri-operative complications and pain after the suburethral sling procedure for urinary stress incontinence: A French prospective randomised multicentre study comparing the retropubic and transobturator routes. *Eur Urol* 2006; **49**:133-138.
44. Minaglia S, Ozel B, Klutke C, Ballard C, Klutke J: Bladder injury during transobturator sling. *Urology* 2004; **64**:376-377.
45. Clemens JQ, DeLancey JO, Faerber GJ, Westney OL, McGuire EJ: Urinary tract erosions after synthetic pubovaginal slings: Diagnosis and management strategy. *Urology* 2000; **56**:589-594.
46. Kobashi KC, Govier FE: Management of vaginal erosion of polypropylene mesh slings. *J Urol* 2003; **169**:2242-2243.
47. Karram MM, Segal JL, Vassallo BJ, Kleeman SD: Complications and untoward effects of the tension-free vaginal tape procedure. *Obstet Gynecol* 2003; **101**:929-932.
48. Huang KH, Kung FT, Liang HM, Chang SY: Management of polypropylene mesh erosion after intravaginal midurethral sling operation for female stress urinary incontinence. *Int Urogynecol J Pelvic Floor Dysfunct* 2005; **16**:437-440.
49. Levin I, Groutz A, Gold R, Pauzner D, Lessing JB, Gordon D: Surgical complications and medium-term outcome results of tension-free vaginal tape: A prospective study of 313 consecutive patients. *Neurourol Urodyn* 2004; **23**:7-9.
50. Charalambous S, Touloupidis S, Fatles G, et al: Transvaginal vs transobturator approach for synthetic sling placement in patients with stress urinary incontinence. *Int Urogynecol J Pelvic Floor Dysfunct* 2007.
51. Siegel AL: Vaginal mesh extrusion associated with use of Mentor transobturator sling. *Urology* 2005; **66**:995-999.
52. Deval B, Ferchaux J, Berry R, et al: Objective and subjective cure rates after trans-obturator tape (OBTAPe) treatment of female urinary incontinence. *Eur Urol* 2006; **49**:373-377.
53. Baessler K, Hewson AD, Tunn R, Schuessler B, Maher CF: Severe mesh complications following intravaginal slingplasty. *Obstet Gynecol* 2005; **106**:713-716.
54. Tsivian A, Kessler O, Mogutin B, et al: Tape related complications of the tension-free vaginal tape procedure. *J Urol* 2004; **171**:762-764.
55. Huang JY, Ziegler C, Tulandi T: Cervical stump necrosis and septic shock after laparoscopic supracervical hysterectomy. *J Minim Invasive Gynecol* 2005; **12**:162-164.
56. Stoeckel W: Über die Verwendung der Muculi Pyramidales bei der Operativen Behandlung der Incontinentia Ulinae. *Zbl Gynals* 1917; **41**:11.
57. Marshall VF, Marchetti AA, Krantz KE: The correction of stress incontinence by simple vesicourethral suspension. 1949. *J Urol* 2002; **167**:1109-1115.
58. Pereyra AJ: A simplified surgical procedure for the correction of stress incontinence in women. *West J Surg Obstet Gynecol* 1959; **67**:223-226.
59. Stamey TA: Endoscopic suspension of the vesical neck for urinary incontinence. *Surg Gynecol Obstet* 1973; **136**:547-554.
60. Raz S: Modified bladder neck suspension for female stress incontinence. *Urology* 1981; **17**:82-85.
61. Ijland MM, Fischer DC, Kieback DG, McGrath G, Farnsworth B: Midline intravaginal slingplasty for treatment of urinary stress incontinence: Results of an independent audit up to 2 years after surgery. *Int Urogynecol J Pelvic Floor Dysfunct* 2005; **16**:447-454.
62. Ulmsten U: An introduction to tension-free vaginal tape (TVT)--A new surgical procedure for treatment of female urinary incontinence. *Int Urogynecol J Pelvic Floor Dysfunct* 2001; **12(Suppl 2)**:S3-S4.
63. Olsson I, Kroon U: A three-year postoperative evaluation of tension-free vaginal tape. *Gynecol Obstet Invest* 1999; **48**:267-269.
64. Wang AC, Lo TS: Tension-free vaginal tape. A minimally invasive solution to stress urinary incontinence in women. *J Reprod Med* 1998; **43**:429-434.
65. Jacquetin B: Use of "TVT" in surgery for female urinary incontinence. *J Gynecol Obstet Biol Reprod (Paris)* 2000; **29**:242-247.
66. Soulie M, Delbert-Julhes F, Cuvillier X, et al: Repair of female urinary incontinence with prolene "TVT": Preliminary results of a multicenter and prospective survey. *Prog Urol* 2000; **10**:622-628.